

## Competition Between Saturated and Unsaturated Components for Reacting with Emerging 1:1 Adduct Radical as Cause for Peaking Dependence of 1:1 Adduct Formation Rate on Unsaturated Component Concentration in Free-Radical Nonbranched-Chain Processes of Initiated Addition to Molecular C=C Bonds in Binary Systems

Michael M. Silaev

Chemistry Faculty, Lomonosov Moscow State University, Vorobievsky Gory, Moscow 119991, Russia.

\*Corresponding author E-mail address: [mmsilaev@rc.chem.msu.ru](mailto:mmsilaev@rc.chem.msu.ru)

ISSN: 2582-3353



### Publication details

Received: 17<sup>th</sup> February 2020  
Revised: 12<sup>th</sup> March 2020  
Accepted: 12<sup>th</sup> March 2020  
Published: 18<sup>th</sup> March 2020

**Abstract:** The reaction scheme is suggested for the initiated nonbranched-chain addition of free radicals to the double bonds of the unsaturated compounds. The proposed scheme includes the reaction competing with chain propagation reactions through a reactive free radical. The chain evolution stage in this scheme involves three types of free radicals. One of them is relatively low-reactive and inhibits the chain process by shortening of the kinetic chain length. Based on the suggested scheme, four rate equations (containing one to three parameters to be determined directly) are deduced using quasi-steady-state treatment. These equations provide good fits for the nonmonotonic (peaking) dependences of the formation rates of the molecular products (1:1 adducts) on the concentration of the unsaturated component in binary systems consisting of a saturated component (hydrocarbon, alcohol, etc.) and an unsaturated component (alkene, allyl alcohol, etc.). The unsaturated compound in these systems is both a reactant and an autoinhibitor generating low-reactive free radicals. It is shown that a maximum in the studied experimental curves arises from the competition between saturated and unsaturated components for reacting with the emerging 1:1 adduct radical.

**Keywords:** Binary System; Unsaturated Compound; Low-Reactive Radical; Autoinhibitor; Competing Reaction; Nonbranched-Chain Addition; Kinetic Equation; Rate; Parameters; Thermochemical Data; Energy

## 1. Introduction

A free radical may be low-reactive if its unpaired  $p$ -electron may be delocalized, e.g., over conjugated bonds as in the case of allyl radical  $\text{CH}_2=\text{CH}\dot{\text{C}}\text{H}_2$ .<sup>1</sup> In nonbranched-chain processes of reactive free radical (addend) addition to double bonds of molecules, the formation of rather low-reactive free radicals in reactions, which are parallel to or competing with propagation via a reactive radicals, lead to chain termination, because these low-reactive radicals (inhibitors) do not participate in further chain propagation and because they decay when colliding with each other or with chain-carrier reactive radicals thus resulting in inefficient expenditure of the latter and process inhibition.

In similar processes involving the addend and inhibitor radicals in diffusion controlled bimolecular chain-termination reactions of three types, the dependences of the rate of molecular 1:1 adduct

formation on the concentration of the unsaturated component (which is the source of low-reactive free radicals in a binary system of saturated and unsaturated components) have a maximum, usually in the region of small (optimal) concentrations.

Here, reactions of addition of reactive free radicals to double bonds of alkene and their derivatives to give 1:1 adduct radicals are taken as examples to consider the role of low-reactive free radicals as inhibitors of the nonbranched chain processes at moderate temperatures.

The 1:1 adduct radical (which is the heaviest and the largest among the free radicals that result from the addition of one addend radical to the double bond of the molecule) may have an increased energy owing to the energy liberated in the transformation of a double bond into an ordinary bond (30–130 kJ mol<sup>-1</sup> for the gas phase under standard conditions).<sup>[1–4]</sup> Therefore, it can decompose or react with one of the surrounding molecules in the place of its formation without diffusing in the solution and, hence, without participating in radical-radical chain termination reactions. Which of the two reactions of the adduct radical, the reaction with the

<sup>1</sup>Note that the activity of a free radical is also connected to the reaction heat in which it participates.

saturated component or the reaction with the unsaturated component, dominates the kinetics of the process will depend on the reactivity and concentration ratios of the components in the binary system.

Earlier,<sup>[5,6]</sup> there were attempts to describe such peaking dependences fragmentarily, assuming that the saturated or unsaturated component is in excess, in terms of the direct and inverse proportionalities, respectively, that result from the simplification of a particular case of the kinetic equation set up by the quasi-steady-state treatment of binary copolymerization involving fairly long chains.<sup>[5]</sup> This specific equation is based on an irrational function, whose plot is a monotonic curve representing the dependence of the product formation rate on the concentration of the unsaturated component. This curve comes out of the origin of coordinates, is convex upward, and has an asymptote parallel to the abscissa axis. Replacing the component concentrations with the corresponding mole fractions generates a peak in this irrational function and thereby makes it suitable to describe the experimental data.<sup>[7]</sup> However, this circumstance cannot serve as a sufficient validation criterion for the mechanism examined, because the new property imparted to the function by the above artificial transformation does not follow from the solution of the set of algebraic equations that are set up for the reaction scheme accepted for the process in a closed system and express the equality of the steady-state formation and disappearance rates of the reactive intermediates.

This publication presents free radical additions to alkenes and their derivatives<sup>[8-11]</sup> yielding various 1:1 molecular adducts, whose formation rates as a function of the unsaturated compound concentration pass through a maximum. In the kinetic description of these non-telomerization chain processes, the reaction between the 1:1 adduct radical and the unsaturated molecule, which is in competition with chain propagation through a reactive free radical ( $\text{PCl}_2$ ,  $\text{C}_2\text{H}_5\dot{\text{C}}\text{HOH}$ , etc.), is included for the first time in the chain propagation stage. This reaction yields a low-reactive radical (such as  $\text{CH}_2=\text{C}(\text{CH}_3)\dot{\text{C}}\text{H}_2$ ) and thus leads to chain termination because this radical does not continue the chain and thereby inhibits the chain process.<sup>[10]</sup> We will consider kinetic variants for the case of comparable component concentrations with an excess of the saturated component<sup>[8,9]</sup> and the case of an overwhelming excess of the saturated component over the unsaturated component<sup>[10,11]</sup> Based on the reaction scheme suggested for the kinetic description of the addition process, we have derived kinetic equations with one to three parameters to be determined directly.

Reducing the number of unknown parameters in a kinetic equation will allow one to decrease the narrowness of the correlation of these parameters and to avoid a sharp build-up of the statistical error in the nonlinear estimation of these parameters in the case of a limited number of experimental data points.<sup>[12]</sup> The rate constant of the addition of a free radical to the double bond of the unsaturated molecule, estimated as a kinetic parameter, can be compared to its reference value if the latter is known. This provides a clear criterion to validate the mathematical description against experimental data.

The kinetic equations were set up using the quasi-steady-state treatment. This method is the most suitable for processes that

include eight to ten or more reactions and four to six different free radicals and are described by curves based on no more than three to seven experimental points. In order to reduce the exponent of the  $2k_5[\text{R}_1^\bullet]^2$  term in the  $d[\text{R}_1^\bullet]/dt = 0$  equation to unity,<sup>[8]</sup> we used the following condition for the early stages of the process:  $k_6 = \sqrt{2k_5 2k_7}$ ,<sup>[16]</sup> and hence,  $V_1 = V_5 + 2V_6 + V_7 = (\sqrt{2k_5} [\text{R}_1^\bullet] + \sqrt{2k_7} [\text{R}_2^\bullet])^2$ . Here,  $[\text{R}_1^\bullet]$  and  $[\text{R}_2^\bullet]$  are the concentrations of the addend radical and the low-reactive (inhibitor) radical, respectively;  $V_1$  is the initiation rate;  $V_5$ ,  $2V_6$ , and  $V_7$  are the rates of the three types of diffusion-controlled quadratic-law chain termination reactions;  $2k_5$  and  $2k_7$  are the rate constants of the loss of identical free radicals via the reactions  $\text{R}_1^\bullet + \text{R}_1^\bullet$  and  $\text{R}_2^\bullet + \text{R}_2^\bullet$ , respectively;  $k_6$  is the rate constant of the loss of different free radicals via the  $\text{R}_1^\bullet + \text{R}_2^\bullet$  reaction (see *Scheme*). The kinetic equations thus obtained fit the peaking rate curves well throughout the range of unsaturated component concentrations in the binary systems. Our mathematical simulation was based on experimental data obtained for  $\gamma$ -radiation-induced addition reactions for which the initiation rate  $V_1$  is known.

The analysis of stable liquid-phase products was carried out by the gas chromatographic method.




## 2. Addition to the C=C Bond of Alkenes and Their Derivatives







When reacting with alkenes not inclined to free-radical polymerization, the free radicals originating from inefficient saturated telogens, such as alcohols<sup>[14]</sup> and amines,<sup>[15]</sup> usually add to the least substituted carbon atom at the double bond, primarily yielding a free 1:1 adduct radical. This radical accumulates an energy of 90–130  $\text{kJ mol}^{-1}$ , which is released upon the transformation of the C=C bond to an ordinary bond (according to the data reported for the addition of nonbranched  $\text{C}_1$ – $\text{C}_4$  alkyl radicals to propene and of similar  $\text{C}_1$  and  $\text{C}_2$  radicals to 1-butene in the gas phase under standard conditions).<sup>[1-4]</sup> Such adduct radicals, which do not decompose readily for structural reasons, can abstract the most labile atom from a neighbour molecule of the saturated or unsaturated component of the binary reaction system, thus turning into a 1:1 adduct molecule. The consecutive and parallel reactions involved in this free-radical nonbranched-chain addition process are presented below (*Scheme*). In the case of comparable component concentrations with a non-overwhelming excess of the saturated component, extra reaction (1b) ( $k_{1b} \neq 0$ ) is included in the initiation stage.<sup>[8,9]</sup> In the case of an overwhelming excess of the saturated component reaction (1b) is ignored ( $k_{1b} = 0$ ).<sup>[10,11]</sup>

### 2.1. Comparable Component Concentrations

#### Scheme

##### Chain initiation



**Chain propagation****Inhibition****Chain termination**

In this scheme, I is an initiator (e.g., a peroxide);<sup>[5]</sup>  $R_0^\bullet$  is a reactive (initiating) radical; A and B are hydrogen or halogen atoms;<sup>[2,5,14-21]</sup>  $R_1^\bullet$  is  $\cdot\text{PCl}_2$ ,<sup>[16]</sup>  $\cdot\text{CCl}_3$ ,<sup>[17]</sup> alkyl,<sup>[2,5]</sup> 1-hydroxyalkyl<sup>[5,6,14,19-21]</sup> or a similar functionalized reactive addend radical;<sup>[5]</sup>  $R_2^\bullet$  is an alkenyl radical (allyl or higher),<sup>[2,5,14-19]</sup> 1-hydroxyalkenyl,<sup>[5,14,15,20,21]</sup> or a similar functionalized low-reactive radical (inhibitor);<sup>[5,15]</sup>  $R_3^\bullet$  is a saturated reactive 1:1 adduct radical;  $R_0A$ ,  $R_0B$ , and  $R_1A$  are saturated molecules;  $R_2B$  is an unsaturated molecule (alkene or its derivative);  $R_3A$  and  $R_3B$  are 1:1 adduct molecules; Prod designates the molecular products resulting from the dimerization or disproportionation of free radicals. The chain evolution (propagation and inhibition) stage of *Scheme* include consecutive reactions 2 and 3, parallel (competing) reaction pairs 3 and 4, and consecutive-parallel reaction pair 2–4.

The initiation reaction 1 is either the decomposition of a chemical initiator<sup>[5,14,15]</sup> or a reaction induced by light<sup>[5,17,18]</sup> or ionizing radiation.<sup>[16-20]</sup> The overall rate of chain initiation (reactions 1, 1a, and 1b) is determined by the rate of the rate-limiting step ( $k_{1b} > k_{1a}$ ). The reaction between the free radical  $R_2^\bullet$ , which results from reactions 1b and 4, and the saturated molecule  $R_1A$  is energetically unfavorable because it implies the formation of the free radical  $R_1^\bullet$ , which is less stable than the initial one. The addition reaction 2 may be accompanied by the abstraction reaction 2a.  $R_1^\bullet + R_2B \xrightarrow{k_{2a}} R_1B + R_2^\bullet$  which yields the product  $R_1B$  via a nonchain mechanism. Reaction 2a does not regenerate the addend radical  $R_1^\bullet$  and is not necessary for a kinetic description of the process, because the rate ratio of reactions 2 and 2a,  $V_2/V_{2a} = k_2/k_{2a}$ , is independent of the concentration of the unsaturated component  $R_2B$  in the system. The inhibition of the nonbranched-chain addition process is due to reaction 4, in which the adduct radical  $R_3^\bullet$  is spent in an inefficient way, since this reaction, unlike reaction 3, does not regenerate  $R_1^\bullet$ . The inhibiting effect is also due to the loss of chain carriers  $R_1^\bullet$  through their collisions with low-reactive unsaturated radicals  $R_2^\bullet$ , but to a much lesser extent.

The rates of the formation ( $V$ , mol dm<sup>-3</sup> s<sup>-1</sup>) of the 1:1 adducts  $R_3A$  (via a chain mechanism) and  $R_3B$  (via a nonchain mechanism) in reactions 3 and 4 are given by the equations

$$V_3(R_3A) = \frac{[\gamma I / (\gamma I + x)] V_1 \alpha k_2 x}{k_2 x^2 + (\alpha I + x) \sqrt{2k_5 V_1}}, \quad (1)$$

$$V_4(R_3B) = \frac{[\gamma I / (\gamma I + x)] V_1 k_2 x^2}{k_2 x^2 + (\alpha I + x) \sqrt{2k_5 V_1}}, \quad (2)$$

where  $V_1$  is the rate of the initiation reaction 1;  $I = [R_1A]$  and  $x = [R_2B]$  are the molar concentrations of the initial components, with  $I > x$ ;  $k_2$  is the rate constant of the addition of the  $R_1^\bullet$  radical from the saturated component  $R_1A$  to the unsaturated molecule  $R_2B$  (reaction 2); and  $\gamma = k_{1a}/k_{1b}$  and  $\alpha = k_3/k_4$  are the rate constant ratios for competing (parallel) reactions ( $\alpha$  is the first chain-transfer constant for the free-radical telomerization process.)<sup>[5]</sup> The rate ratio for the competing reactions is  $V_3/V_4 = \alpha I/x$ , and the chain length is  $\nu = V_3/V_1$ . Earlier mathematical simulation<sup>[8]</sup> demonstrated that replacing the adduct radical  $R_3$  with the radical  $R_2$ <sup>[5]</sup> in the reaction between identical radicals and in the reaction involving  $R_1$  gives rise to a peak in the curve of the 1:1 adduct formation rate as a function of the concentration of the unsaturated component. Reaction 1b, which is in competition with reaction 1a, is responsible for the maximum in the curve described by Eq. (2), and reaction 4, which is in competition with reaction (3), is responsible for the maximum in the curve defined by Eq. (1).


The number of unknown kinetic parameters to be determined directly ( $k_2$ ,  $\alpha$ , and  $\gamma$ ) can be reduced by introducing the condition  $\gamma \cong \alpha$ , which is suggested by the chemical analogy between the competing reactions pairs 1a–1b and 3–4. For example, the ratios of the rate constants of the reactions of  $\cdot\text{OH}$ ,  $\text{CH}_3\text{O}^\bullet$ ,  $\cdot\text{CH}_3$ ,  $\text{NO}_3^\bullet$ , and  $\text{H}_2\text{PO}_4^\bullet$  with methanol to the rate constants of the reactions of the same radicals with ethanol in aqueous solution at room temperature are 0.4–0.5.<sup>[22,23]</sup> For the same purpose, the rate constant of reaction 2 in the kinetic equation can be replaced with its analytical expression  $k_2 = \alpha I_m \sqrt{2k_5 V_1} / x_m^2$ , which is obtained by solving the quadratic equation following from the reaction rate extremum condition  $\partial V_{3,4}(1:1 \text{ Adduct}) / \partial x = 0$ , where  $V_{3,4}(1:1 \text{ Adduct}) = V_3 + V_4$ . After these transformations, the overall formation rate equation for the 1:1 adducts  $R_3A$  and  $R_3B$  (which may be identical, as in the case of  $R_3H$ <sup>[5,8,9,15-18]</sup>) appears as

$$V_{3,4}(1:1 \text{ Adduct}) = \frac{V_1 \alpha k_2 x}{k_2 x^2 + (\alpha I + x) \sqrt{2k_5 V_1}} = \quad (3)$$

$$= \frac{V_1 \alpha I x}{x^2 + (\alpha I + x) x_m^2 / \alpha I_m}, \quad (3a)$$

where  $I_m$  and  $x_m$  are the component concentrations  $I$  and  $x$  at the points of maximum of the function. Provided that  $V_1$  is known, the only parameter in Eq. (3a) to be determined directly is  $\alpha$ . If  $V_1$  is known only for the saturated component  $R_1A$ , then, for the binary system containing comparable  $R_1A$  and  $R_2B$  concentrations, it is better to use the quantity  $\lambda V_1$ , where  $\lambda = I/(I + x)$  is the mole fraction of  $R_1A$ , in place of  $V_1$  in Eqs. (3) and (3a).

The two variable concentrations in the kinetic equation (3) –  $I$  and  $x$  – can be reduced to one variable by replacing them with the



**Fig. 1.** Reconstruction of the functional dependences (curves) of the product formation rates  $V_{3,4}$  (1,  $\nabla$ ) on the mole fraction of the unsaturated component ( $\chi$ ) from empirical data (symbols) using Eq. (3b) (model optimization with respect to the parameter  $\alpha$ ) for the phosphorus trichloride–methylpropene reaction system at 303 K<sup>[16]</sup> (standard deviation of  $S_V = 2.58 \times 10^{-6}$ ) and (2,  $\circ$ ) on the concentration of the unsaturated component ( $x$ ) from empirical data (symbols) using Eq. (4a) (model optimization with respect to  $V_1$ ,  $x_m$ , and  $\alpha$ ) for the 2-propanol–2-propen-1-ol system at 433 K<sup>[20]</sup> ( $S_V = 5.91 \times 10^{-7}$ ).

corresponding mole fractions. Substituting the expression  $k_2 = \left\{ \alpha \left[ \left( \frac{1}{x_m} \right) - 1 \right]^2 - 1 \right\} \sqrt{2k_5 V_1} / (l_m + x_m)$ , derived from the rate extremum condition, into this transformed equation for the binary system containing comparable component concentrations, we obtain

$$V_{3,4}(\text{Adduct 1:1}) = \frac{V_1 \alpha (1 - \chi) \chi}{\chi^2 + [\alpha(1 - \chi) + \chi] \left\{ \alpha \left[ \left( \frac{1}{x_m} \right) - 1 \right]^2 - 1 \right\}}, \quad (3b)$$

where  $1 - \chi = l / (l + x)$  and  $\chi = x / (l + x)$  are the mole fractions of the components  $R_1A$  and  $R_2B$  ( $0 < \chi < 1$ ), respectively, and  $x_m$  is the  $\chi$  value at the point of maximum.

The overall formation rate of the 1:1 adducts  $R_3A$  and  $R_3B$  is a sophisticated function of the formation and disappearance rates of the radicals  $R_1^\bullet$  and  $R_2^\bullet$ :  $V(R_3A, R_3B) = (V_{1a} + V_3 - V_5) - (V_{1b} + V_4 - V_7)$ .

The application of the above rate equations to particular single nonbranched-chain additions is illustrated in Fig. 1. Curve 1 represents the results of simulation in terms of Eq. (3b) for the observed 1:1 adduct formation rate as a function of the mole fraction of the unsaturated component in the phosphorus trichloride–methylpropene<sup>2</sup> reaction system at 303 K.<sup>[16]</sup> In this simulation, the <sup>60</sup>Co  $\gamma$ -radiation dose rate was set at  $P = 0.01 \text{ Gy s}^{-1}$  and the initiation yield was taken to be  $G(^{\bullet}PCl_2) = 2.8$  particles per 100 eV ( $1.60 \times 10^{-17} \text{ J}$ ) of the energy absorbed by the solution.<sup>[16]</sup> The product of reaction 3 is  $Cl_2PCH_2C(Cl)(CH_3)CH_3$  (two isomers),  $V_1 = 4.65 \times 10^{-9} \text{ mol dm}^{-3} \text{ s}^{-1}$  at  $\chi = 0$ , and  $2k_5 = 3.2 \times 10^8 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ . This leads to  $\alpha = (2.5 \pm 0.4) \times 10^3$ , and the rate constant of reaction 2 derived from this  $\alpha$  value is

$$k_2 = (1.1 \pm 0.2) \times 10^4 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}.$$

## 2.2. Excess of the Saturated Component

If the concentration of the saturated component exceeds the concentration of the unsaturated component in the binary system, reaction 1b can be neglected. If this is the case ( $k_{1b} = 0$ ), then, in the numerators of the rate equations for reactions 3 and 4 (Eqs. (1) and (2)),  $\eta / (\eta + x) = 1$  and the overall rate equation for the formation of the 1:1 adducts  $R_3A$  and  $R_3B$  will appear as

$$V_{3+4}(\text{1:1 Adduct}) = \frac{V_1(\alpha l + x)k_2 x}{k_2 x^2 + (\alpha l + x)\sqrt{2k_5 V_1}} = \quad (4)$$

$$= \frac{V_1 x}{\frac{x^2}{\alpha l + x} + \left( \frac{\sqrt{\alpha l_m}}{x_m} + \frac{1}{\sqrt{\alpha l_m}} \right)^{-2}} \quad (4a)$$

where the parameters are designated in the same way as in Eqs. (1)–(3a),  $l \gg x$ , and  $k_2 = \left[ \left( \frac{\sqrt{\alpha l_m}}{x_m} \right) + \left( \frac{1}{\sqrt{\alpha l_m}} \right) \right]^2 \sqrt{2k_5 V_1}$  is determined from the condition  $\partial V_{3,4}(\text{1:1 Adduct}) / \partial x = 0$ .

The rate equations for the chain termination steps (reactions 5–7 in the *Scheme*;  $k_{1b} = 0$ ) are identical to the equations reported for these steps in<sup>[24]</sup> if  $\beta = 0$  in those equations.

Note that, if it is necessary to supplement *Scheme* for  $k_{1b} = 0$  with the formation of  $R_1B$  via the possible nonchain reaction 2a (which is considered in the Section 2.1), the parameter  $k_{2a}$  should be included in the denominator of Eq. (4) to obtain  $k_2 x^2 + (\alpha l + x)(k_{2a} x + \sqrt{2k_5 V_1})$ .

The analytical expression for  $k_2$  in the case of  $k_{2a} \neq 0$  is identical to the expression for  $k_2$  for Eq. (4). The equation for the rate  $V_{2a}(R_1B)$  can be derived by replacing  $k_2$  with  $k_{2a}$  in the numerator of Eq. (4) containing  $k_{2a}$  in its denominator.

For approximate estimation of the parameters of the kinetic equations (3) and (4), Eq. (4) under the conditions (a)  $k_2 x^2 \ll (\alpha l + x)\sqrt{2k_5 V_1}$  (ascending branch of a peaked curve) and (b)  $k_2 x^2 \gg (\alpha l + x)\sqrt{2k_5 V_1}$  (descending branch) is transformed into simple functions (direct and inverse proportionality, respectively) of the concentration  $x$  of the unsaturated compound, which allow tentative estimates of the parameters  $k_2$  and  $\alpha$  to be derived from the experimental 1:1 adduct formation rate  $V_{3+4}$  provided that  $V_1$  and  $2k_5$  are known:

$$V_{3+4} = \sqrt{V_1} k_2 x / \varphi \sqrt{2k_5} \quad (5)$$

$$V_{3+4} = (V_1 / \varphi) [(\alpha l / x) + 1] \quad (6)$$

where  $\varphi = 1$  under conditions (a) and (b) and  $\varphi = 2$  at the point of maximum (where  $k_2 x^2 \cong (\alpha l + x)\sqrt{2k_5 V_1}$ ).

The equation (3) presented in the previous section reduces into Eq. (5) if the condition (a) is fulfilled and  $\alpha l \gg x$  (in this case, at the beginning of an ascending branch,  $\varphi = 1$  only) and into Eq. (7), which express a simple inversely proportional dependence of the reaction rate  $V_{3,4}$  on  $x$  and provide tentative estimates of the parameter  $\alpha$ :

<sup>2</sup>In an earlier work,<sup>[10]</sup> the methylpropene concentration in this system was overvalued by a factor of 1.7 when it was derived from the mole fractions given in.<sup>[19]</sup>

$$V_{3,4} = V_1 \alpha I / \varphi x \quad (7)$$

where  $\varphi = 2$  at the point of maximum (where  $k_2 x^2 \cong (\alpha I + x) \sqrt{2k_5 V_1}$ ) and  $\varphi = 1$  for the descending branch of the curve.

Curve 2 in Fig. 1 illustrates the good fit between Eq. (4a) and the observed 1:1 adduct formation rate as a function of the concentration of the unsaturated component in the reaction system 2-propanol-2-propen-1-ol at 433 K.<sup>[8,9]</sup> In this description, we used a <sup>60</sup>Co  $\gamma$ -radiation dose rate of  $P = 4.47 \text{ Gy s}^{-1}$ .<sup>[20]</sup> The product of reactions 3 and 4 is  $\text{CH}_3(\text{CH}_3)\text{C}(\text{OH})\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ , and  $2k_5 = 1.0 \times 10^{10} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ . The following parameters were obtained:  $V_1 = (3.18 \pm 0.4) \times 10^6 \text{ mol dm}^{-3} \text{ s}^{-1}$ ,  $x_m = (3.9 \pm 0.5) \times 10^{-2} \text{ mol dm}^{-3}$ , and  $\alpha = (6.8 \pm 0.8) \times 10^{-2}$ . The rate constant of reaction 2 derived from this  $\alpha$  is  $k_2 = (1.0 \pm 0.14) \times 10^5 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ .

### 3. Conclusions

In summary, the material on the kinetics of nonbranched-chain addition of free saturated radicals to double bonds of molecules of alkene and its derivative makes it possible to describe, using rate equations (3) and (4) obtained by quasi-steady-state treatment, experimental dependences with a maximum of the formation rates of molecular 1:1 adducts on the concentration of an unsaturated compound over the entire region of its change in binary reaction systems consisting of saturated and unsaturated components (Fig. 1).

The proposed addition mechanism involves the reaction of a free 1:1 adduct radical with an unsaturated molecule yielding a low-reactive free radical (the reaction 4 competing with the chain propagation reaction 3 in Scheme). In such reaction systems, the unsaturated compound is both a reactant and an autoinhibitor, specifically a source of low-reactive free radicals shortening kinetic chains.

The kinetic equations obtained on the basis of the suggested reaction scheme involving two competitive reactions 3 and 4 at the chain evolution stage make it possible to directly determine the rate constant  $k_2$  of the addition reaction 2 without using a separate labor-intensive method of competing reactions as often happens in such cases.

The optimum concentration  $x_m$  of unsaturated component in the binary system at which the process rate is maximal can be derived with the help of obtained kinetic equations (3a) and (4a) or from the corresponding analytical expressions for  $k_2$  if other parameters are known. This opens a way to intensification of some technological processes that are based on the addition of free radicals to the double bonds of unsaturated molecules and occur via a nonbranched-chain mechanism through the formation of 1:1 adducts.

### Conflicts of Interest

The authors declare no conflict of interest.



© 2020, by the authors. Licensee Ariviyal Publishing, India. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

### References

- Gurvich L.V.; Karachevtsev G.V.; Kondratev V.N.; Yu. A. Lebedev; Medvedev V.A.; Potapov V. K.; Yu. S. Khodeev. Bond Dissociation Energies, Ionization Potentials, and Electron Affinity. Nauka, Moscow, 1974, 80-91.
- Benson S.W. Thermochemical Kinetics: Methods for the Estimation of Thermochemical Data and Rate Parameters. 2<sup>nd</sup> Edition, Wiley, New York, 1976. [\[Link\]](#)
- Pedley J.B.; Naylor R.D.; Kirby S.P. Thermochemical Data of Organic Compounds. 2<sup>nd</sup> Edition, Chapman & Hall, London, 1986. [\[Link\]](#)
- Yu. D. Orlov; Yu. A. Lebedev; I. Sh. Saifullin. Thermochemistry of Organic Free Radicals. Nauka, Moscow, 2001.
- Ch. Walling. Free Radicals in Solution. Wiley, New York, 1956.
- Emanuel N.M.; Denisov E.T.; Maizus Z.K. Chain Oxidation Reactions of Hydrocarbons in the Liquid Phase. Nauka, Moscow, 1965.
- Poluektov V.A.; Babkina E.I.; Begishev I.R. On the Dependence of the Rate of a Chain Reaction on the Reactant Ratio. *Dokady Akademii Nauk SSSR*, 1974, **215**, 649–652.
- Silaev M.M. Competition Kinetics of Nonbranched Chain Processes of Free-Radical Addition to Double Bonds of Molecules with the Formation of 1:1 Adducts. *Kinet. Catal.*, 1999, **40**, 256–259. [\[Link\]](#)
- Silaev M.M. Simulation of the Nonbranched-Chain Addition of Saturated Free Radicals to Alkenes and Their Derivatives Yielding 1:1 Adducts. *Theor. Found. Chem. Eng.*, 2007, **41**, 273–278. [\[Link\]](#)
- Silaev M.M.; Bugaenko L.T. Mathematical Simulation of the Kinetics of Radiation Induced Hydroxyalkylation of Aliphatic Saturated Alcohols. *Radiat. Phys. Chem.*, 1992, **40**, 1–10. [\[Link\]](#)
- Silaev M.M.; Bugaenko L.T. Kinetics of the Addition of  $\alpha$ -Hydroxyalkyl Radicals to 2-Propen-1-ol and Formaldehyde. *Kinet. Catal.*, 1994, **35**, 509–513.
- Bard Y. Nonlinear Parameter Estimation. Academic, New York, 1974. [\[Link\]](#)
- Bateman L. Alkene Oxidation. *Quarterly Reviews*, 1954, **8**, 147–167.
- Urry W.H.; Stacey F.W.; Huyser E.S.; Juveland O.O. The Peroxide- and Light-Induced Additions of Alcohols to Alkenes. *J. Am. Chem. Soc.*, 1954, **76**, 450–455. [\[Link\]](#)
- Urry W.H.; Juveland O.O. Free Radical Additions of Amines to Alkenes. *J. Am. Chem. Soc.*, 1958, **80**, 3322–3328. [\[Link\]](#)
- Shostenko A.G.; Zagorets P.A.; Dodonov A.M.; Greish A.A.  $\gamma$ -Radiation-Induced Addition of Phosphorus Trichloride to Isobutylene. *Khimiya Vysokikh Energii*, 1970, **4**, 357.
- Kim V.; Shostenko A.G.; Gasparyan M.D. Reactivity of Polychloroalkyl Radicals in the Telomerization of  $\text{CCl}_4$  with 1-Propene and 2-Methyl-1-Propene. *React. Kinet. Catal. Lett.*, 1979, **12**, 479–484. [\[Link\]](#)
- Myshkin V.E.; Shostenko A.G.; Zagorets P.A.; Markova K.G.; Pchelkin A.I. Determination of Absolute Rate Constants for the Addition of the Ethyl Radical to Alkenes. *Teoreticheskaya i Eksperimental'naya Khimiya*, 1977, **13**, 266–271.
- Zamyslov R.A.; Shostenko A.G.; Dobrov I.V.; Tarasova N.P. Kinetics of  $\gamma$ -Radiation-Induced Reactions of 2-Propanol with Trifluoropropene and Hexafluoro-propene. *Kinetika i Katalyz.*, 1987, **28**, 977–979.
- Silaev M.M. Dependence of Radiation-Chemical  $\gamma$ -Diol Yields on the 2-Propen-1-ol Concentration in the Radiolysis of Aliphatic Saturated  $\text{C}_1$ – $\text{C}_3$  Alcohol-2-Propen-1-ol Systems. *High Energy Chem.*, 1990, **24**, 245–246. [\[Link\]](#)
- Silaev M.M.  $\gamma$ -Diol Formation via the Autooxidation of 2-Propen-1-ol Solutions in Saturated Alcohols. *Vestnik Moskovskogo Universiteta, Ser. 2: Khimiya*, 1994, **35**, 40–42.
- Bugaenko L.T.; Kuzmin M.G.; Polak L.S. *High-Energy Chemistry*. Horwood Hall, New York, 1993, 112.
- Thomas J.K. Pulse Radiolysis of Aqueous Solutions of Methyl Iodide and Methyl Bromide. The Reactions of Iodine Atoms and Methyl Radicals in Water. *J. Phys. Chem.*, 1967, **71**, 1919–1925. [\[CrossRef\]](#)
- Silaev M.M. The Competition Kinetics of Radical-Chain Addition. *Russ. J. Phys. Chem.*, 1999, **73**, 1050–1054. [\[Link\]](#)